

RIGIDITY THEOREMS FOR LAGRANGIAN SUBMANIFOLDS OF C^n AND CP^n WITH CONFORMAL MASLOV FORM

XIAOLI CHAO AND YUXIN DONG*

ABSTRACT. In this paper, we obtain a rigidity theorem for Lagrangian submanifolds of C^n and CP^n with conformal Maslov form.

1. Introduction

Let $(\widetilde{M}^n, J, g)$ be a Kähler manifold of complex dimension n . The Kähler form ω on $\widetilde{M}^n$ is given by $\omega(X, Y) = g(X, JY)$. An immersion $\psi : M^n \rightarrow \widetilde{M}^n$ of an n -dimensional manifold M is called Lagrangian if $\psi^*\omega \equiv 0$. Lagrangian submanifolds in a Kähler manifold or more generally in a symplectic manifold appear naturally in the context of mathematical physics. Since 1970s, various kind of Lagrangian submanifolds in Kähler manifolds have also been investigated extensively from Riemannian geometric point of view (see [1] and the references therein).

From [4], we know that the n -sphere cannot be embedded in C^n as a Lagrangian submanifold. This result is not true when the Lagrangian sphere is immersed but not embedded. The simplest immersions of S^n into C^n , known as Whitney spheres [8], are induced by a map $\psi : E^{n+1} \rightarrow C^n (\cong R^{2n})$ defined by

$$\psi(x_0, x_1, \dots, x_n) = \frac{r}{1 + x_0^2}(x_1, \dots, x_n, x_0x_1, \dots, x_0x_n) + A$$

where r is a positive number and A is a vector in C^n . We will refer to r and A as the radius and the center of the Whitney sphere respectively. The Whitney spheres have the best possible behavior either from the viewpoint of topology or the viewpoint of submanifold geometry. They have a unique self-intersection point $\psi(-1, 0, \dots, 0) = \psi(1, 0, \dots, 0)$. Their second fundamental form h satisfy

$$(1) \quad h(X, Y) = \frac{n}{n+2} \{ \langle X, Y \rangle H + \langle JX, H \rangle JY + \langle JY, H \rangle JX \}$$

where X, Y are tangent to ψ and $H = \frac{1}{n} \text{trace}(h)$ is the mean curvature vector of the Whitney sphere. This property may be regarded as the Lagrangian version of umbilicity. In [7], it was proved that the Whitney spheres are the only closed Lagrangian submanifolds in C^n having this property.

1991 *Mathematics Subject Classification.* 53C40, 53C42.

*Supported by Zhongdian grant of NSFC

From [1], we know that there is no Lagrangian immersion of S^n into C^n with parallel mean curvature vector. However, the Whitney spheres have the property that JH are conformal vector fields. We may regard a Lagrangian submanifold with this property as the analogue of hypersurfaces of constant mean curvature. As the dual form of JH is the Maslov form of the Lagrangian immersion, the Lagrangian submanifolds whose JH are conformal vector field will be known as Lagrangian submanifolds with conformal Maslov form (see [7]). In [7], these submanifolds were studied when the ambient space is C^n . The authors proved in [7] that the Whitney spheres are the only compact Lagrangian submanifolds of C^n with conformal Maslov form and the null first Betti number. This result may be regarded as the Lagrangian version of the classical Hopf theorem. When the ambient spaces are CP^n and CH^n , the authors in [2], [4] constructed similar Lagrangian immersions from S^n into these spaces, which were called Whitney spheres too. It turns out that (1) also characterizes Whitney spheres among closed non-minimal Lagrangian submanifolds when the ambient spaces are C^n and CP^n . In the case of CH^n , there are other two families of examples, besides the Whitney spheres of CH^n , that satisfy (1).

In this paper, we first introduce a modified second fundamental form B suggested by (1). In terms of the square norm of B , we will establish rigidity theorems for Lagrangian submanifolds of C^n and CP^n with conformal Maslov form, and thus characterize the Whitney spheres (Theorem 3.3) in these spaces.

2. Preliminaries

Let $\psi : M^n \rightarrow \widetilde{M}^n$ be a Lagrangian immersion of an n -dimensional manifold M^n into a Kähler manifold $\widetilde{M}^n$. Let $\nabla, \widetilde{\nabla}$ denote the Levi-Civita connections of M and $\widetilde{M}$ respectively. The second fundamental form h of ψ is given by

$$h(X, Y) = \widetilde{\nabla}_X Y - \nabla_X Y$$

for any vector fields X, Y tangent to M . The mean curvature vector H of ψ is defined by

$$H = \frac{1}{n} \text{trace}(h) = \sum_{i=1}^n h(e_i, e_i)$$

where $\{e_1, \dots, e_n\}$ is an orthonormal tangent frame of M .

We choose a local orthonormal frame field $\{e_1, \dots, e_n\}$ of M and hence $\{e_{1^*} = J e_1, \dots, e_{n^*} = J e_n\}$ forms a local normal frame field of M in $\widetilde{M}^n$. Let

$$\{\omega^1, \dots, \omega^n, \omega^{1^*}, \dots, \omega^{n^*}\}$$

denote the dual frame field of $\{e_1, \dots, e_n, e_{1^*}, \dots, e_{n^*}\}$.

In this paper, we shall make use of the following convention of indices:

$$1 \leq A, B, C, \dots \leq 2n, \quad 1 \leq i, j, k, m, l \dots \leq n$$

The structure equations of $\widetilde{M}$ are given by

$$\begin{aligned} d\omega^A &= - \sum_B \omega_B^A \wedge \omega^B, \quad \omega_B^A + \omega_A^B = 0 \\ d\omega_B^A &= - \sum_C \omega_C^A \wedge \omega_B^C + \Omega_B^A, \quad \Omega_B^A = \sum_{CD} \frac{1}{2} K_{BCD}^A \omega^C \wedge \omega^D \\ K_{BCD}^A + K_{BDC}^A &= 0 \end{aligned}$$

Restricting these forms to M , we have

$$\begin{aligned}
\omega_j^i + \omega_i^j &= 0, \quad \omega_j^i = \omega_{j*}^{i*}, \quad \omega_j^{i*} = \omega_i^{j*}, \\
\omega_i^{m*} &= h_{ij}^{m*} \omega^j, \quad h_{ij}^{m*} = h_{ji}^{m*} = h_{mj}^{i*} = h_{im}^{j*}, \\
d\omega^i &= -\omega_j^i \wedge \omega^j, \quad \omega_j^i + \omega_i^j = 0 \\
d\omega_j^i &= -\omega_k^i \wedge \omega_j^k + \frac{1}{2} R_{jkl}^i \omega^k \wedge \omega^l, \quad R_{jkl}^i + R_{jlk}^i = 0 \\
R_{jkl}^i &= K_{jkl}^i + \sum_m (h_{jl}^{m*} h_{ik}^{m*} - h_{il}^{m*} h_{jk}^{m*}) \\
R_{j*kl}^{i*} &= K_{j*kl}^{i*} + \sum_m (h_{lm}^{j*} h_{mk}^{i*} - h_{ml}^{i*} h_{mk}^{j*})
\end{aligned}$$

Define the first covariant derivative of h_{ij}^{m*} by

$$\sum_k h_{ijk}^{m*} \omega^k = dh_{ij}^{m*} - \sum_k h_{kj}^{m*} \omega_i^k - \sum_k h_{ik}^{m*} \omega_j^k + \sum_k h_{ij}^{k*} \omega_k^{m*}$$

We denote by $\widetilde{M}^n(4c)$ the complex space form of constant holomorphic sectional curvature $4c$, i.e., its curvature satisfies

$$K_{BCD}^A = c\{\delta_{AC}\delta_{BD} - \delta_{AD}\delta_{BC} + J_{AC}J_{BD} - J_{AD}J_{BC} + 2J_{AB}J_{CD}\}$$

where

$$(J_{AB}) = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}$$

It is known that

$$\widetilde{M}^n(4c) \cong \begin{cases} C^n, & \text{if } c = 0 \\ CP^n, & \text{if } c > 0 \\ CH^n, & \text{if } c < 0 \end{cases}$$

For a Lagrangian submanifold M in $\widetilde{M}^n(4c)$, we obtain the Codazzi equation by a direct computation

$$h_{ijk}^{m*} = h_{ikj}^{m*}$$

Notice that JH is a tangent vector field when the submanifold is Lagrangian.

Lemma 2.1. ([7]) *JH is a conformal vector field if and only if*

$$\nabla JH = \text{div}(JH/n)I,$$

which is equivalent to

$$\sum_k h_{kk,l}^{m*} = -\text{div}(JH)\delta_l^m$$

Definition 2.2. A Lagrangian submanifold M is said to have conformal Maslov form if JH is a conformal vector field of M .

In the introduction, we present the Whitney spheres in C^n . From [4], we obtain the following family of Lagrangian immersions

$$\psi_\theta : S^n \rightarrow CP^n, \quad \theta > 0$$

given by

$$\psi_\theta(x_1, \dots, x_n, x_{n+1}) = \left[\left(\frac{(x_1, \dots, x_n)}{ch_\theta + ish_\theta x_{n+1}}, \frac{sh_\theta ch_\theta (1 + x_{n+1}^2) + ix_{n+1}}{ch_\theta^2 + sh_\theta^2 x_{n+1}^2} \right) \right]$$

that are called the Whitney spheres in CP^n . There are Whitney immersions of S^n into CH^n too (see [4] for details).

Lemma 2.3. ([1], [4], [7]) *Let $\psi : M \rightarrow \widetilde{M}^n(4c)$ be a Lagrangian immersion of an n -dimensional manifold M in the complex space form $\widetilde{M}^n(4c)$ with $c \geq 0$. Then the second fundamental form h of ψ satisfies*

$$h(X, Y) = \frac{n}{n+2} \{ \langle X, Y \rangle H + \langle JX, H \rangle JY + \langle JY, H \rangle JX \}$$

for any vectors X and Y tangent to M if and only if either M is a totally geodesic submanifold or it is an open set of the Whitney Sphere.

Remark 2.1. In the case $c < 0$, there are other two families of examples, besides Whitney spheres of CH^n , whose second fundamental forms satisfy the property in Lemma 2.3.

3. Rigidity theorems for Lagrangian submanifolds

In this section, we will establish a rigidity theorem for Lagrangian submanifolds in C^n and CP^n with conformal Maslov form.

Let $\psi : M^n \rightarrow \widetilde{M}^n(4c)$ be a Lagrangian submanifold with conformal Maslov form. We define

$$(2) \quad B(X, Y) = h(X, Y) - \frac{n}{n+2} \{ \langle X, Y \rangle H + \langle JX, H \rangle JY + \langle JY, H \rangle JX \}$$

for any tangent vector fields X, Y of M . It is easy to verify that $\text{trace}(B) = 0$. With respect to the above orthonormal frame field $\{e_1, \dots, e_n, e_{1^*}, \dots, e_{n^*}\}$ along M , we have

$$\begin{aligned} h_{ij}^{m*} &= b_{ij}^{m*} + \frac{n}{n+2} \{ H^{m*} \delta_{ij} + \langle H, J e_i \rangle \delta_{jm} + \langle H, J e_j \rangle \delta_{im} \} \\ &= b_{ij}^{m*} + \frac{n}{n+2} \{ H^{m*} \delta_{ij} + H^{i*} \delta_{jm} + H^{j*} \delta_{im} \} \\ &= b_{ij}^{m*} + c_{ij}^{m*} \end{aligned}$$

where

$$c_{ij}^{m*} = \frac{n}{n+2} \{ H^{m*} \delta_{ij} + H^{i*} \delta_{jm} + H^{j*} \delta_{im} \}$$

By a direct computation, we may get

$$(3) \quad \|B\|^2 = \|h\|^2 - \frac{3n^2}{n+2}|H|^2$$

Define the first and the second covariant derivatives of b_{ij}^{m*} as follows:

$$\begin{aligned} \sum_k b_{ijk}^{m*} \omega^k &= db_{ij}^{m*} - \sum_k b_{kj}^{m*} \omega_i^k - \sum_k b_{ik}^{m*} \omega_j^k + \sum_k b_{ij}^{k*} \omega_k^{m*} \\ \sum_l b_{ijkl}^{m*} \omega^l &= db_{ijk}^{m*} - \sum_l b_{ljk}^{m*} \omega_i^l - \sum_l b_{ilk}^{m*} \omega_j^l - \sum_l b_{ijl}^{m*} \omega_k^l + \sum_l b_{ijk}^{l*} \omega_l^{m*} \end{aligned}$$

Lemma 3.1. *The tensor field B is a Codazzi tensor field, i.e., $b_{ijk}^{m*} = b_{ikj}^{m*}$.*

Proof. For any point of M , we may take a local orthonormal frame $\{e_i\}_{i=1}^n$ near p such that $(\nabla e_i)_p = 0$ for $i = 1, \dots, n$, which imply that $(\nabla^\perp e_i^*)_p = 0$ for $i = 1, \dots, n$. By using Lemma 2.1, we have

$$\begin{aligned} b_{ijk}^{m*} &= h_{ijk}^{m*} - \frac{n}{n+2} \{H_{,k}^{m*} \delta_{ij} + \langle H_{,k}, Je_i \rangle \langle Je_j, e_m^* \rangle + \langle H_{,k}, Je_j \rangle \langle Je_i, e_m^* \rangle\} \\ &= h_{ijk}^{m*} - \frac{n}{n+2} \{H_{,k}^{m*} \delta_{ij} + H_{,k}^{i*} \delta_{jm} + H_{,k}^{j*} \delta_{im}\} \\ &= h_{ijk}^{m*} + \frac{\text{div}(JH)}{n+2} \{\delta_{km} \delta_{ij} + \delta_{ik} \delta_{jm} + \delta_{jk} \delta_{im}\} \end{aligned}$$

Therefore $b_{ijk}^{m*} = b_{ikj}^{m*}$. $\square$

The Laplacian of b_{ij}^{m*} is defined by

$$(4) \quad \Delta b_{ij}^{m*} = \sum_k b_{ijkk}^{m*} = \sum_k b_{ikjk}^{m*} = \sum_k b_{kijk}^{m*}$$

By Ricci identities and Lemma 3.1, we obtain

$$\begin{aligned} b_{kijk}^{m*} &= b_{kikj}^{m*} + \sum_t b_{kt}^{m*} R_{ijk}^t + \sum_t b_{ti}^{m*} R_{kjk}^t - \sum_t b_{ki}^{t*} R_{t*jk}^{m*} \\ (5) \quad &= b_{kkij}^{m*} + \sum_t b_{kt}^{m*} R_{ijk}^t + \sum_t b_{ti}^{m*} R_{kjk}^t - \sum_t b_{ki}^{t*} R_{t*jk}^{m*} \end{aligned}$$

Now we compute

$$\frac{1}{2} \Delta \|B\|^2 = \sum_{ijkm} (b_{ijk}^{m*})^2 + \sum_{ijm} b_{ij}^{m*} \Delta b_{ij}^{m*}$$

From (4),(5), we have

$$\begin{aligned} \sum_{ijm} b_{ij}^{m*} \Delta b_{ij}^{m*} &= \underbrace{\sum_{ijkmt} b_{ij}^{m*} b_{kt}^{m*} R_{ijk}^t}_{(I)} + \underbrace{\sum_{ijkmt} b_{ij}^{m*} b_{ti}^{m*} R_{kjk}^t}_{(II)} - \underbrace{\sum_{ijkmt} b_{ij}^{m*} b_{ki}^{t*} R_{t*jk}^{m*}}_{(III)} \end{aligned}$$

where

$$\begin{aligned}
(I) &= c \sum_{ijkmt} b_{ij}^{m*} b_{kt}^{m*} (\delta_{tj} \delta_{ik} - \delta_{tk} \delta_{ij}) + \sum_{ijkmtl} b_{ij}^{m*} b_{kt}^{m*} (b_{tj}^{l*} b_{ik}^{l*} - b_{tk}^{l*} b_{ij}^{l*}) \\
&+ \sum_{ijkmtl} b_{ij}^{m*} b_{kt}^{m*} \{b_{tj}^{l*} c_{ik}^{l*} + c_{tj}^{l*} b_{ik}^{l*} + c_{tj}^{l*} c_{ik}^{l*} - b_{tk}^{l*} c_{ij}^{l*} - c_{tk}^{l*} c_{ij}^{l*}\} \\
&= c \|B\|^2 + \sum_{ijkmtl} b_{ij}^{m*} b_{kt}^{m*} (b_{tj}^{l*} b_{ik}^{l*} - b_{tk}^{l*} b_{ij}^{l*}) + \frac{2n}{n+2} \sum_{jklmt} b_{jk}^{m*} b_{kt}^{m*} b_{tj}^{l*} H^{l*} \\
&+ \sum_{ijkmtl} b_{ij}^{m*} b_{kt}^{m*} (c_{tj}^{l*} c_{ik}^{l*} - c_{tk}^{l*} c_{ij}^{l*}) \\
&= c \|B\|^2 + \frac{n^2}{(n+2)^2} \|B\|^2 |H|^2 + \sum_{ijkmtl} b_{ij}^{m*} b_{kt}^{m*} (b_{tj}^{l*} b_{ik}^{l*} - b_{tk}^{l*} b_{ij}^{l*}) \\
&+ \frac{2n}{n+2} \sum_{jklmt} b_{jk}^{m*} b_{kt}^{m*} b_{tj}^{l*} H^{l*} + \frac{2n^2}{(n+2)^2} \sum_{ijkmt} b_{ij}^{m*} b_{jk}^{m*} H^{i*} H^{k*}
\end{aligned}$$

$$\begin{aligned}
(II) &= (n-1)c \|B\|^2 + \sum_{ijklmt} b_{ij}^{m*} b_{ti}^{m*} \{n b_{tj}^{l*} H^{l*} + n c_{tj}^{l*} H^{l*} - b_{tk}^{l*} b_{kj}^{l*} - b_{tk}^{l*} c_{kj}^{l*} \\
&- c_{tk}^{l*} b_{kj}^{l*} - c_{tk}^{l*} c_{kj}^{l*}\} \\
&= (n-1)c \|B\|^2 - \sum_{ijklmt} b_{ij}^{m*} b_{ti}^{m*} b_{tk}^{l*} b_{kj}^{l*} + \frac{n^2 - 2n}{n+2} \sum_{ijlmt} b_{ij}^{m*} b_{ti}^{m*} b_{tj}^{l*} H^{l*} \\
&+ \frac{n^2}{n+2} \{ \|B\|^2 |H|^2 + 2 \sum_{ijmt} b_{ij}^{m*} b_{ti}^{m*} H^{j*} H^{t*} \} - \sum_{ijklmt} b_{ij}^{m*} b_{ti}^{m*} c_{tk}^{l*} c_{kj}^{l*} \\
&= (n-1)c \|B\|^2 + \frac{n^3}{(n+2)^2} \|B\|^2 |H|^2 - \sum_{ijkmtl} b_{ij}^{m*} b_{ti}^{m*} b_{tk}^{l*} b_{kj}^{l*} \\
&+ \frac{n^2 - 2n}{n+2} \sum_{ijlmt} b_{ij}^{m*} b_{ti}^{m*} b_{tj}^{l*} H^{l*} + \frac{n^2(n-2)}{(n+2)^2} \sum_{ijmt} b_{ij}^{m*} b_{ti}^{m*} H^{j*} H^{t*}
\end{aligned}$$

$$\begin{aligned}
(III) &= \sum_{ijklmt} b_{ij}^{m*} b_{ki}^{t*} \{b_{lj}^{m*} b_{lk}^{t*} - b_{lk}^{m*} b_{lj}^{t*} + b_{lj}^{m*} c_{lk}^{t*} + c_{lj}^{m*} b_{lk}^{t*} - b_{lk}^{m*} c_{lj}^{t*} - c_{lk}^{m*} b_{lj}^{t*} \\
&+ c_{lj}^{m*} c_{lk}^{t*} - c_{lk}^{m*} c_{lj}^{t*}\} - c \|B\|^2 \\
&= -c \|B\|^2 - \frac{n^2}{(n+2)^2} \|B\|^2 |H|^2 + \sum_{ijklmt} b_{ij}^{m*} b_{ki}^{t*} (b_{lj}^{m*} b_{lk}^{t*} - b_{lk}^{m*} b_{lj}^{t*}) \\
&- \frac{2n^2}{(n+2)^2} \sum_{ijkmt} b_{ij}^{m*} b_{jk}^{m*} H^{i*} H^{k*} - \frac{2n}{n+2} \sum_{ijlmt} b_{ij}^{m*} b_{lj}^{m*} b_{li}^{t*} H^{t*}
\end{aligned}$$

Set $B_{m^*} = (b_{ij}^{m^*})$. Then we get

$$\begin{aligned}
\frac{1}{2}\Delta\|B\|^2 &= \sum_{ijk m} (b_{ijk}^{m^*})^2 + \sum_{ijk m} b_{ij}^{m^*} b_{ijk}^{m^*} \\
&= \sum_{ijk m} (b_{ijk}^{m^*})^2 + (n+1)c\|B\|^2 + \frac{n^2}{(n+2)}\|B\|^2|H|^2 \\
&\quad + \sum_{ijk m t l} b_{ij}^{m^*} b_{kt}^{m^*} (b_{tj}^{l^*} b_{ik}^{l^*} - b_{tk}^{l^*} b_{ij}^{l^*}) - \sum_{ijk m t l} b_{ij}^{m^*} b_{ki}^{t^*} (b_{lj}^{m^*} b_{lk}^{t^*} - b_{lk}^{m^*} b_{lj}^{t^*}) \\
&\quad - \sum_{ijk m t l} b_{ij}^{m^*} b_{it}^{m^*} b_{tk}^{l^*} b_{kj}^{l^*} + n \sum_{ij l m t} b_{ij}^{m^*} b_{jl}^{m^*} b_{li}^{t^*} H^{t^*} + \frac{n^2}{n+2} \sum_{ijk m} b_{ij}^{m^*} b_{jk}^{m^*} H^{i^*} H^{k^*} \\
&= \sum_{ijk m} (b_{ijk}^{m^*})^2 + (n+1)c\|B\|^2 + \frac{n^2}{(n+2)}\|B\|^2|H|^2 + \sum_{ij} \text{tr}(B_{i^*} B_{j^*} - B_{j^*} B_{i^*})^2 \\
&\quad - \sum_{ij} (\text{tr} B_{i^*} B_{j^*})^2 + \frac{n^2}{n+2} \sum_{ijk m} b_{ij}^{m^*} b_{jk}^{m^*} H^{i^*} H^{k^*} + n \sum_{ij m t} b_{ij}^{m^*} b_{jl}^{m^*} b_{li}^{t^*} H^{t^*} \\
&\geq \sum_{ijk m} (b_{ijk}^{m^*})^2 + (n+1)c\|B\|^2 + \frac{n^2}{(n+2)}\|B\|^2|H|^2 - \sum_i S_{i^*} - 2 \sum_{i \neq j} S_{i^*} S_{j^*} \\
&\quad + n|H| \sum_i S_{i^*} \lambda_i + \frac{n^2}{n+2} S_{1^*} |H|
\end{aligned}$$

where

$$S_{i^*} = \sum_{jt} (b_{jt}^{i^*})^2, \quad b_{ij}^{1^*} = \lambda_i \delta_{ij}, \quad S_H = \sum_i \lambda_i^2$$

Here we assume $e_{1^*} // H$ and $S_H = \sum (b_{ij}^{1^*})^2$.

For a matrix $A = (a_{ij})$, we denote by $N(A)$ the square norm of A as in [3], i.e.,

$$N(A) = \text{trace}(A^t A) = \sum_{ij} (a_{ij})^2$$

Lemma 3.2. ([6]) Let $A_1, \dots, A_p$ be symmetric $(n \times n)$ -matrices ($p \geq 2$). Denote $S_{\alpha\beta} = \text{trace}(A_\alpha A_\beta)$, $S_\alpha = S_{\alpha\alpha} = N(A_\alpha)$, $S = \sum_{i=1}^p S_i$. Then

$$\sum_{\alpha\beta} N(A_\alpha A_\beta - A_\beta A_\alpha) + \sum_{\alpha\beta} S_{\alpha\beta}^2 \leq \frac{3}{2} S^2$$

From Lemma 3.2, we know

$$\begin{aligned}
\frac{1}{2}\Delta\|B\|^2 &\geq \sum_{ijkm} (b_{ijk}^{m*})^2 + (n+1)c\|B\|^2 + \frac{n^2}{(n+2)}\|B\|^2|H|^2 - \frac{n+2}{2}\sum_i S_{i*}^2 \\
&\quad - 2\sum_{i\neq j} S_{i*}S_{j*} + \frac{n}{2}\sum_i (\lambda_i^2|H|^2 + 2|H|S_{i*}\lambda_i + S_{i*}^2) \\
&\geq (n+1)c\|B\|^2 + \frac{n^2}{(n+2)}\|B\|^2|H|^2 - \frac{n+2}{2}(\sum_i S_{i*}^2 + 2\sum_{i\neq j} S_{i*}S_{j*}) \\
&\quad + n\sum_{i\neq j} S_{i*}S_{j*} + \frac{n}{2}\sum_i (\lambda_i|H| + S_{i*})^2 \\
&\geq (n+1)c\|B\|^2 + \frac{n^2}{(n+2)}\|B\|^2|H|^2 - \frac{3(n+2)}{4}\|B\|^4
\end{aligned}$$

Theorem 3.3. *Let $\psi : M \longrightarrow \widetilde{M}^n(4c)$ ($n \geq 2$) be a (non-minimal) Lagrangian immersion of an n -dimensional closed manifold M into the complex space form $\widetilde{M}^n(4c)$ ($c \geq 0$) with conformal Maslov form. If*

$$\|B\|^2 \leq \frac{4(n+1)c}{3(n+2)} + \frac{4n^2|H|^2}{3(n+2)^2}$$

or equivalently,

$$\|h\|^2 \leq \frac{4(n+1)c}{3(n+2)} + \frac{n^2(9n+22)|H|^2}{3(n+2)^2}$$

then $\psi(M)$ is the Whitney sphere.

Proof. Since M is closed, we have

$$\begin{aligned}
0 &= \int_M \frac{1}{2}\Delta\|B\|^2 \geq \int_M \left\{ (n+1)c\|B\|^2 + \frac{n^2}{(n+2)}\|B\|^2|H|^2 - \frac{3(n+2)}{4}\|B\|^4 \right\} \\
&= \int_M \|B\|^2 \left\{ (n+1)c + \frac{n^2}{(n+2)}|H|^2 - \frac{3(n+2)}{4}\|B\|^2 \right\}
\end{aligned}$$

By the assumption

$$\|B\|^2 \leq \frac{4(n+1)c}{3(n+2)} + \frac{4n^2|H|^2}{3(n+2)^2}$$

thus we get $B = 0$ on M , i.e.,

$$h(X, Y) = \frac{n}{n+2} \{g(X, Y)H + g(JX, H)JY + g(JY, H)JX\}$$

for any tangent vector fields X, Y of M . It follows from Lemma 2.3 that $\psi(M)$ is the Whitney sphere. From (3), we see that the condition for $\|B\|^2$ is equivalent to

$$\|h\|^2 \leq \frac{4(n+1)c}{3(n+2)} + \frac{n^2(9n+22)|H|^2}{3(n+2)^2}$$

This complete the proof of the Theorem. $\square$

Remark 3.1. (a) The non-minimality condition is void for the case $c = 0$. Obviously (3) tells us that ([7],[2])

$$\|h\|^2 \geq \frac{3n^2|H|^2}{n+2}$$

(b) Let $M = S^1(r_1) \times \cdots \times S^1(r_n) \hookrightarrow C^1 \times \cdots \times C^1 = C^n$ be the standard Lagrangian flat torus in C^n , which has parallel mean curvature vector. By a direct computation, we know that it has conformal Maslov form and

$$\|B\|^2 = \frac{n^2(n-1)H^2}{n+2}$$

We also have the standard flat Lagrangian torus in $\widetilde{M}^n(4c)$ with $c > 0$ (see for example [Ch]), which has parallel mean curvature, and thus has conformal Maslov form. These examples show that some upper bound for $\|B\|^2$ is necessary to characterize Whitney spheres. It would be interesting to get the optimal gap theorem.

Corollary 3.4. *Let $\psi : M \longrightarrow \widetilde{M}^n(4c)$ ($n \geq 2$) be a (non-minimal) Lagrangian immersion of an n -dimensional closed manifold M into complex space form $\widetilde{M}^n(4c)$ ($c > 0$) with conformal Maslov form. If*

$$\|B\|^2 \leq \frac{4(n+1)c}{3(n+2)}$$

then $\psi(M)$ is the Whitney sphere.

REFERENCES

- [1] B. Y. Chen, *Riemannian geometry of Lagrangian submanifolds*, Taiwanese J. of Math., **5**(2001), 681-723.
- [2] B. Y. Chen, *Jacobi's elliptic functions and Lagrangian immersions*, Proc. Royal Soc. Edin., **126** (1996), 687-704.
- [3] S. S. Chern, M. do Carmo and S. Kobayashi, *Minimal submanifolds of a sphere with second fundamental form of constant length*, Functional Analysis and Related Fields (F. E. Browder ed.), Springer-Verlag, New York (1970).
- [4] I. Castro, C.R. Montealegre and F. Urbano, *Closed conformal vector fields and Lagrangian submanifolds in complex space forms*, Pacific J. Math., **199** (2001), 269-301.
- [5] M. L. Gromov, *Pseudoholomorphic curves in symplectic manifolds*, invent. Math., **82** (1985), 307-347.
- [6] A.M. Li and J.M. Li, *An intrinsic theorem for minimal submanifolds in a sphere*, Archiv der Math., **58** (1992), 582-594.
- [7] A. Ros and F. Urbano, *Lagrangian submanifolds of C^n with conformal Maslov form and the Whitney sphere*, J. Math. Soc. Japan, **50** (1998), 203-226.
- [8] A. Weinstein, *Lectures on symplectic manifolds*, Conference board of the Mathematical Scientific, **29**(1977).

Xiaoli Chao
Department of Mathematics,
Southeast University, Nanjing, 210096,
P. R. China
Email address: xlchao@seu.edu.cn

and

Yuxin Dong
Institute of Mathematics,
Fudan University, Shanghai, 200433,
P. R. China
And
Key Laboratory of Mathematics for Nonlinear Sciences (Fudan University),
Ministry Education
Email address: yxdong@fudan.edu.cn